

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

Product Information

Description: The ChromoTek Ubiquitin-Trap Magnetic Agarose Kit consists of an anti-Ubiquitin Nanobody/VHH, which is coupled to magnetic agarose beads. It also contains lysis, wash, and elution buffers that can be used for the immunoprecipitation of ubiquitin and ubiquitinated proteins from the cell extracts of various species including human, mouse, hamster, dog, yeast, and plants.

Applications: IP, Co-IP

Specificity/Target: Binds to monomeric ubiquitin, ubiquitin chains, and ubiquitinated proteins. Ubiquitin chains may be linked via lysines; linkages via lysines such as K11, K48 and K63 are compatible with binding by the Ubiquitin-Trap (further linkage types also likely to be compatible, but not tested).

Bead Size: 40 µm (cross-linked 6 % magnetic agarose beads)

Wash Buffer Compatibility: 2M NaCl, 5 mM DTT, 0 mM β-mercaptoethanol, 5 mM TCEP, 2% NP40, 2% Triton X-100, 0% SDS, 2-3 M Urea

Type: Nanobody

Class: Recombinant

Host: Camelid

Shipment: Shipped at ambient temperature

Storage Buffer: 20 % ethanol


Storage Condition: Upon receipt store at +4°C. Do not freeze!

Stability: Stable for 1 year upon receipt

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

Selected Validation Data



Immunoprecipitation of Ubiquitin from HEK293T cell extracts (treated with MG-132) using Ubiquitin-Trap Magnetic Agarose Kit.

Kit Components

Component	Composition	Quantity
Ubiquitin-Trap Magnetic Agarose	Anti-ubiquitin VHH cross-linked with magnetic agarose beads	20 rxns (500 ul slurry)
Lysis Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5 % Nonidet™ P40 Substitute, 0.09 % sodium azide	30 mL
RIPA Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.1 % SDS, 1 % Triton™ X-100, 1 % Deoxycholate, 0.09 % sodium azide	30 ml
Dilution Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.018 % sodium azide	50 mL (after dilution with 40 mL water)
Wash Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.05 % Nonidet™ P40 Substitute, 0.5 mM EDTA, 0.018 % sodium azide	50 mL (after dilution with 40 mL water)
Acidic elution buffer	200 mM glycine pH 2.5	3 x 1 mL

Note: Use your equivalent cell lysis buffer for other cell types like yeast, plants, insects, bacteria. Consider using a Wash buffer without detergent for Co-IP.

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

Required Buffer Solutions

Buffer	Composition
2x SDS-sample buffer	120 mM Tris/Cl pH 6.8, 20% glycerol, 4% SDS, 0.04% bromophenol blue, 10% β-mercaptoethanol
Neutralization buffer	1M Tris pH 10.4 (adjust the pH at +4°C

Product Sizes

Product	Product Code	Size
Ubiquitin-Trap Magnetic Agarose Kit	utmak-20	20 reactions including buffers

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

Protocol at a glance

General

- Perform all steps at 4°C
- Use your preferred cell lysis buffer and cell lysis conditions

Cell Lysis




- Use 10^6 - 10^7 cells and 200 μ L Lysis buffer.
- Perform cell lysis and clear lysate
- Mix 200 μ L cleared lysate with 300 μ L dilution buffer.

Bead Equilibration



- Transfer 25 μ L bead slurry into a 1.5 mL tube
- Equilibrate beads 3x with 500 μ L dilution buffer

Protein binding



- Add 500 μ L diluted lysate to beads
- Rotate end-over-end for 1 hour at 4°C.

Washing



- Wash beads 3x with 500 μ L wash buffer
- Transfer beads to a new tube during the last washing step

Elution with SDS-sample buffer



- Resuspend beads in 80 μ L 2x SDS-sample buffer
- Boil beads for 5 min at 95°C
- Analyze the supernatant in SDS-PAGE/ Western Blot

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

Immunoprecipitation Protocol

Cell Material

The following protocol describes the preparation of a mammalian cell lysate.

For other type of cells, we recommend using 500 µg of cell extract and start the protocol with step Bead equilibration.

Note: Precipitation of ubiquitinated proteins may be facilitated by treating cells with proteasome inhibitors prior to harvesting. A commonly used inhibitor is MG-132 (also known as MG132 or carbobenzoxy-Leu-Leu-leucinal), for example. Exact conditions must be optimized for each cell type etc.; a good starting point is to incubate cells with f.c. 5-25 µM MG-132 for 1-2 h. Over-exposure to MG-132 can lead to cytotoxic effects, however.

Mammalian Cell Lysis

Note: Harvesting of cells and cell lysis should be performed with ice-cold buffers. We strongly recommend adding protease inhibitors to the Lysis buffer to prevent degradation of your target protein and its binding partners.

Note: If necessary, deubiquitylates (DUBs), i.e. ubiquitin hydrolases, may be inhibited by the addition of either EDTA or EGTA and either f.c. 5-10 mM iodoacetamide (IAA) or N-ethylmaleimide (NEM). Concentrations may need to be optimized for each cell type, growth conditions etc.

For one immunoprecipitation reaction, we recommend using $\sim 10^6$ - 10^7 cells.

1. Choice of lysis buffer:
 - a. For cytoplasmic proteins, resuspend the cell pellet in 200 µL ice-cold Lysis buffer by pipetting up and down. Supplement Lysis buffer with protease inhibitor cocktail and 1 mM PMSF (not included).
 - b. For nuclear/chromatin proteins, resuspend cell pellet in 200 µL ice-cold RIPA buffer supplemented with DNase I (f.c. 75-150 Kunitz U/mL), MgCl₂ (f.c. 2.5 mM), protease inhibitor cocktail and PMSF (f.c. 1 mM) (not included).
2. Place the tube on ice for 30 min and extensively pipette the suspension every 10 min.
3. Centrifuge cell lysate at 17,000x g for 10 min at +4°C. Transfer cleared lysate (supernatant) to a pre-cooled tube and add 300 µL Dilution buffer supplemented with 1 mM PMSF and protease inhibitor cocktail (not included). If required, save 50 µL of diluted lysate for further analysis (input fraction).

Bead Equilibration

1. Resuspend the beads by gently pipetting them up and down or by inverting the tube. Do not vortex the beads!
2. Transfer 25 µL of bead slurry into a 1.5 mL reaction tube.
3. Add 500 µL ice-cold Dilution buffer.

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

4. Separate the beads with a magnet until the supernatant is clear. Discard the supernatant.

Protein Binding

1. Add diluted lysate to the equilibrated beads.
2. Rotate end-over-end for 1 hour at +4°C.

Washing

1. Separate the beads with a magnet until the supernatant is clear.
2. If required, save 50 µL of supernatant for further analysis (flow-through/non-bound fraction).
3. Discard remaining supernatant.
4. Resuspend beads in 500 µL Wash buffer.
5. Separate the beads with a magnet until the supernatant is clear. Discard the remaining supernatant.
6. Repeat this step at least twice.
7. During the last washing step, transfer the beads to a new tube.

Optional: To increase stringency of the Wash buffer, test various salt concentrations e.g. 150-500 mM, and/or add a non-ionic detergent e.g. Triton™ X-100 (see Wash buffer compatibility information for maximal concentrations).

Elution with 2x SDS-sample buffer (Laemmli)

1. Remove the remaining supernatant.
2. Resuspend beads in 80 µL 2x SDS-sample buffer.
3. Boil beads for 5 min at +95°C to dissociate immunocomplexes from beads.
4. Separate the beads with a magnet.
5. Analyze the supernatant in SDS-PAGE / Western Blot.

Note: For Western blot detection we recommend Ubiquitin Recombinant antibody (80992-1-RR) or Ubiquitin Polyclonal antibody (10201-2-AP) in combination with HRP-conjugated Affinipure Goat Anti-Rabbit IgG (H+L) (SA00001-2)

Elution with Acidic Elution Buffer

1. Remove the remaining supernatant.
2. Add 50-100 µL Acidic elution buffer and constantly pipette up and down for 30-60 sec at +4°C or room temperature.
3. Separate the beads with a magnet until the supernatant is clear.
4. Transfer the supernatant to a new tube.
5. Immediately neutralize the eluate fraction with 5-10 µL Neutralization buffer.
6. Repeat this step at least once to increase elution efficiency.

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

Note: Elution at room temperature is more efficient than elution at +4°C. Prewarm buffers for elution at room temperature.

Related Products

Product	Code
Ubiquitin-Trap Agarose	uta
Ubiquitin-Trap Agarose Kit	utak
Ubiquitin-Trap Magnetic Agarose	utma

Contact

Proteintech North America (HQ)

Proteintech Group, Inc.
5500 Pearl Street, Suite 400
Rosemont, IL 60018 USA

1-888-472-4522
proteintech@ptglab.com

Proteintech Europe

Transmission (6th Fl)
6 Atherton Street
M3 3GS, Manchester, UK

+44 161 839 3007
europe@ptglab.com

ChromoTek & Proteintech Germany

Am Klopferspitz 19
82152, Planegg-Martinsreid
Germany

+49 89 124 148 850
germany@ptglab.com

chromotek® Ubiquitin-Trap Magnetic Agarose Kit

Product Code: utmak

Disclaimer

For research use only (RUO), not for diagnostic or therapeutic use.

ChromoTek® is a registered trademark of ChromoTek GmbH, part of Proteintech Group.